

Table 2. Plant-Back Restrictions for Forage Cover Crops after Post-Emergence Herbicide Applications in Corn

All numbers represent months. A zero indicates no replant restrictions following herbicide application. If a box for a herbicide-species combination has a dash, the restriction is too long to allow for planting of that forage cover crop species within the same growing season as the corn crop.

Herbicide	Cool-Season Species											Warm-Season Species				
	Annual Ryegrass	Barley	Cereal Rye	Clover	Field Pea	Oats	Radish	Triticale	Turnip	Vetch	Wheat	Chickpea	Cowpea	Pearl Millet	Proso Millet	Sorghum
Aim	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Armezon	3	3	3	—	—	3	—	3	—	—	3	—	—	3	3	—
Basis Blend	—	3	3	—	—	3	—	3	—	—	3	—	—	3	3	—
Beacon ^a	—	3	3	—	—	—	—	—	—	—	3	—	—	—	—	—
Buctril ^b	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Cadet	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Callisto	0	—	—	—	—	0	—	—	—	—	—	—	—	0	—	0
Callisto GT	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0
Clarity ^c	3	3	—	—	—	3	—	3	—	—	3	—	—	3	3	—
Dicamba ^d	3	3	—	—	—	3	—	3	—	—	3	—	—	3	3	0.5
DiFlexx ^e	—	2	—	—	—	2	—	—	—	—	2	—	—	—	—	2
Expert	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0
Glyphosate	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Halex GT	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0
Harmony SG	1.5	0	1.5	1.5	1.5	0	1.5	0	1.5	1.5	0	1.5	1.5	1.5	1.5	1.5
Impact	—	3	3	—	—	3	—	—	—	—	3	—	—	—	—	—
Liberty 280	—	2.3	2.3	—	—	2.3	2.3	2.3	2.3	—	2.3	—	—	—	—	—
Permit	2	2	2	—	—	2	—	2	—	—	2	—	—	2	2	2
Require Q ^f	—	—	—	—	—	—	—	—	—	—	3	—	—	—	—	—
Resolve Q	—	—	—	—	—	—	—	—	—	—	3	—	—	—	—	—
Resource	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Solida ^g	—	—	—	—	—	—	—	—	—	—	3	—	—	—	—	—
Solstice	—	—	—	—	—	0	—	—	—	—	—	—	—	0	—	0
Spirit	—	3	3	—	—	3	—	3	—	—	3	—	—	—	—	—
Status ^h	—	1	1	—	—	1	—	1	—	—	1	—	—	1	1	1
Warrant ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0
Yukon	2	2	2	—	—	2	—	2	—	—	2	—	—	2	2	2

See footnotes on next page.

-
- ^a Injury may occur if dry weather prevails during much of the time between Beacon application and seeding of winter cereals or other crops.
 - ^b Do not cut crop for feed, fodder, or graze within 45 days of application.
 - ^c Sorghum may be planted 15 days after Clarity application of 8 oz/ac; all other intervals are for rates of 24 oz/ac or less. See label for rotation intervals at rates of more than 24 oz/ac.
 - ^d Sorghum may be planted 15 days after Dicamba application of 8 oz/ac; all other intervals are for rates of 24 oz/ac or less. See label for rotation intervals at rates of more than 24 oz/ac.
 - ^e When applied at a rate of 24 oz/ac or less. See label for rotation intervals at rates of more than 24 oz/ac.
 - ^f When applied at a rate of 4 oz/ac.
 - ^g When applied at a rate of 1 oz/ac.
 - ^h If at least 1 inch of rainfall or overhead irrigation has been received following the last application of Status (less than or equal to 5 oz/ac only), alfalfa, cereal grain crops, and sorghum may be planted 30 days after the rainfall/irrigation event.
 - ⁱ When planting sorghum, use only seed properly treated with seed protectant or safener.