



Nebraska On-Farm Research Network

UNL Pre-Plant Corn Nitrogen Recommendation vs. Producer Nitrogen Recommendation

*Protocol developed by: Dr. Charles Wortmann, UNL Extension Nutrient Management Specialist
Gary Zoubek and Keith Glewen, UNL Extension Educators*

Objective: Verify the UNL N recommendation by comparing it to the producer's current N recommendation strategy.

Rationale: UNL research consistently confirms that when N is considered to be non-limiting to crop growth, each addition of N fertilizer results in less yield gain. This results in lower N use efficiency and lower net return. This is attributed, at least partly, to the high N use efficiency possible with crops that have healthy and well-developed root systems efficient in nutrient recovery and have plants that are efficient in converting nutrients and carbohydrates to yield (i.e. internal or physiological efficiency). The UNL N recommendation has been **validated** for maximizing profitability for high yielding environments (i.e. >240 bu/ac), as well as lower yield environments over diverse production situations. However, many producers feel that N in excess of the recommendations is needed; this results in reduced profitability and more N loss to the environment.

Procedure: The two treatments in these trials are:

Treatment 1: UNL N recommendation

Treatment 2: Producer's N recommendation

The UNL nitrogen rate can be determined using the following [spreadsheet](#) (use the tabs at the bottom of the spreadsheet for help and instructions). (If link does not work visit the www.cropwatch.unl.edu/farmresearch and click on the research protocols page.) This trial does not require a guidance system or yield mapping capabilities but these are preferred. Residual nitrate-nitrogen sample is required for the UNL N recommendation.

Treatment Design: With two treatments for this trial we are able to use the paired comparison design below. A total of 7 replications should be implemented and harvested. The same hybrid and management practices (other than N) should be used across the entire study area.

*****NOTE!** When designing a nitrogen comparison you need to remember nitrogen is a mobile nutrient and corn roots will spread laterally (i.e. corn plants can take up N from up to a row away). Therefore, the width of the treatment strips shown below must take this into account and compensate for it. We compensate for this by allowing for a "buffer" between the different treatments. This "buffer" area is not used for the yield comparison. For example, if you have a 16 row nitrogen applicator and an 8 row corn head, you will need to harvest the center 8 rows of each 16 row treatment, leaving 4 rows on each side of each treatment strip as "buffer". This area will be harvested but not included in the yields for statistical analysis of the treatments. The following layout demonstrates a 16 row applicator and 8 row corn head. If you have different equipment sizes you will need to adjust accordingly. If you have any questions about the treatment design when working with N, please contact the Nebraska On-Farm Research Network.

Replication

N Application

Harvest

Rep 1	(16 rows) Treatment 1: UNL N recommendation	←4 rows buffer
		← Record Yield from center 8 rows
	(16 rows) Treatment 2: Producer's N recommendation	←4 rows buffer
		←4 rows buffer
Rep 2	(16 rows) Treatment 2: Producer's N recommendation	←4 rows buffer
		← Record Yield from center 8 rows
	(16 rows) Treatment 1: UNL N recommendation	←4 rows buffer
		←4 rows buffer
Rep 3	(16 rows) Treatment 1: UNL N recommendation	←4 rows buffer
		← Record Yield from center 8 rows
	(16 rows) Treatment 2: Producer's N recommendation	←4 rows buffer
		←4 rows buffer
Rep 4	(16 rows) Treatment 2: Producer's N recommendation	←4 rows buffer
		← Record Yield from center 8 rows
	(16 rows) Treatment 1: UNL N recommendation	←4 rows buffer
		←4 rows buffer
Rep 5	(16 rows) Treatment 1: UNL N recommendation	←4 rows buffer
		← Record Yield from center 8 rows
	(16 rows) Treatment 2: Producer's N recommendation	←4 rows buffer
		←4 rows buffer
Rep 6	(16 rows) Treatment 2: Producer's N recommendation	←4 rows buffer
		← Record Yield from center 8 rows
	(16 rows) Treatment 1: UNL N recommendation	←4 rows buffer
		←4 rows buffer
Rep 7	(16 rows) Treatment 1: UNL N recommendation	←4 rows buffer
		← Record Yield from center 8 rows
	(16 rows) Treatment 2: Producer's N recommendation	←4 rows buffer
		←4 rows buffer

Grower Requirements:

1. Flag or **mark** GPS location of each treatment.
2. Provide all necessary **inputs** for crop production.
3. Complete a **background** agronomic form about site and practices.
4. Collect **yield data** and **grain moisture** with weigh wagon or yield monitor. If using yield monitor, please designate a separate “load” for each treatment and set up separate “products” names for each treatment harvested. Yield monitor must be **well calibrated**. Contact UNL Extension if assistance with this process is needed.
5. Collect stand counts at harvest. Each treatment in all replications should have a stand count recorded. It is recommended that at least 3 counts be averaged together for each reported stand count.
6. Submit harvest data to UNL Extension within 30 days of harvest or by Dec. 15 of the harvest year.
7. Allow UNL Extension to use submitted and collected data for research, educational, and informational purposes.

Nebraska On-Farm Research Network will:

1. Provide technical assistance in setting up replicated and randomized experimental design.
2. Provide assistance upon request with treatment implementation, flagging, stand counts, stalk rot tests, and recording yield.
3. Analyze raw data using statistical analysis and provide this information to the grower.

For assistance with studies, please contact the Nebraska On-Farm Research Network Coordinators:

Keith Glewen: kglewen1@unl.edu or 402-624-8005

Laura Thompson: laura.thompson@unl.edu or 402-624-8033

Or your local educator

Disclaimer: The Nebraska On-Farm Research Network does not endorse the use of products tested in on-farm replicated strip trials. While treatments are replicated within trials and may be replicated across multiple sites under various conditions, your individual results may vary.

Copyright ©2015

Extension is a Division of the Institute of Agriculture and Natural Resources at the University of Nebraska–Lincoln cooperating with the Counties and the United States Department of Agriculture.

University of Nebraska–Lincoln Extension educational programs abide with the nondiscrimination policies of the University of Nebraska–Lincoln and the United States Department of Agriculture.

