

**Table 4. Plant-Back Restrictions for Forage Cover Crops after Post-Emergence Herbicide Applications in Soybeans**

All numbers represent months. A zero indicates no replant restrictions following herbicide application. If a box for a herbicide-species combination has a dash, the restriction is too long to allow for planting of that forage cover crop species within the same growing season as the corn crop.

Herbicide	Cool-Season Species											Warm-Season Species			
	Annual Ryegrass	Barley	Cereal Rye	Clover	Field Pea	Oats	Radish	Triticale	Turnip	Vetch	Wheat	Chickpea	Cowpea	Pearl Millet	Sorghum
Assure II	—	—	—	—	0	—	—	—	—	—	—	0	0	—	—
Cadet	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Classic <sup>1</sup>	3	3	3	—	—	3	—	—	—	—	3	—	—	—	—
Cobra	2	2	2	—	—	2	—	2	—	—	2	—	—	2	2
Extreme	—	—	—	—	0	—	—	—	—	—	—	—	0	—	—
Fusilade DX	—	2	2	—	—	—	—	2	—	—	2	—	—	2	2
Glyphosate	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Harmony SG	1.5	0	1.5	1.5	1.5	0	1.5	0	1.5	1.5	0	1.5	1.5	1.5	1.5
Liberty	—	70 days	70 days	—	—	70 days	70 days	70 days	70 days	—	70 days	—	—	70 days	70 days
Poast Plus	1	1	1	0	1	1	1	1	—	1	—	—	1	1	1
Pursuit	—	—	—	—	0	—	—	—	—	—	—	—	0	—	—
Raptor	—	—	—	—	0	—	—	—	—	—	3	—	0	—	—
Resource	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Select Max	1	1	1	0	0	1	0	1	0	1	1	1	0	1	1
Synchrony	3	3	3	—	—	3	—	3	—	—	3	—	—	3	—
Targa	—	0	—	—	0	—	—	—	—	—	0	—	0	—	—
Torment	—	—	—	—	—	—	—	—	—	—	—	0	—	—	—
Ultra Blazer <sup>2</sup>	100 days	40 days	40 days	100 days	100 days	40 days	100 days	40 days	100 days	100 days	40 days	100 days	100 days	40 days	100 days

<sup>1</sup> Classic should be applied early post-emergence to meet the three-month interval for planting listed forage cover crops in fall.

<sup>2</sup> Application rate should not be more than 1.5 pint/ac. Soybean cannot be used for livestock grazing or feeding after application of Ultra Blazer.