## Table 3. Plant-Back Restrictions for Forage Cover Crops after Pre-Emergence Herbicide Applications in Soybeans

All numbers represent months. A zero indicates no replant restrictions following herbicide application. If a box for a herbicide-species combination has a dash, the restriction is too long to allow for planting of that forage cover crop species within the same growing season as the corn crop.

	Cool-Season Species												Warm-Season Species				
Herbicide	Annual Ryegrass	Barley	Cereal Rye	Clover	Field Pea	Oats	Radish	Triticale	Turnip	Vetch	Wheat	Chickpea	Cowpea	Pearl Millet	Proso Millet	Sorghum	
Anthem <sup>a</sup>		_	_	_	_	4	_	_	_	_	4	_	_	_	_	_	
Anthem Maxx <sup>b</sup>	_	_	_	_	_	4	_	_	_	_	4	_	_	_	_	_	
Authority Assist <sup>c</sup>	_	_	4	_	_		_	_	_	_	4	_	_	-	—	_	
Authority Elite	_	4.5	4.5	_	_	0	4.5	_	_	_	4.5	_	_	_	_	_	
Authority First	_	_	_	_	—	_	_	—	—	_	4	_	—	_	—	—	
Authority Maxx	_	4	4	_	_	_	_	_	_	_	4	_	_	-	—	_	
Authority MTZ	_	4	4	_	—	_	_	—	—	_	4	_	—	_	—	_	
Authority XL	_	4	_	_	—	_	_	—	—	_	4	_	—	_	—	_	
Boundary	_	4.5	_	_	_	_	_	_	_	_	4.5	_	_	-	—	_	
Canopy	4	4	4	_	—	_	_	—	—	_	4	_	—	_	—	_	
Command <sup>d</sup>	_	-		I	_	0/9 <sup>e</sup>	_	_	_	_	_	_	_	_	—	_	
Dimetric DF	_	4	_	_	—	_	_	—	—	_	4	_	—	_	—	_	
Dual II Magnum <sup>f</sup>	4.5	4.5	4.5	I	_		_	_	2	_	4.5	2	_	4.5	—	2	
Enlite	_	4	4	_	_	_	—	—	_	_	3	_	—		_	_	
Envive	4	4	4	_	—	_	_	—	—	_	4	_	—	_	—	_	
FirstRate	_	-		I	_		_	_	_	_	4	_	_	_	—	_	
Gangster	_	_	_	_	—	_	_	—	—	_	3	_	—	_	—	_	
Optill	_	_	4	_	—	4	_	—	—	_	4	_	4	_	—	_	
Optill PRO	_	_	4	_	_	4	—	—	_	_	4	_	—		_	_	
Outlook	_	4	4	I	_		_	_	_	_	4	_	_	4	—	_	
Prefix	_	4.5	4.5	_	_	_	_	_	_	_	4.5	_	_	4.5	_	_	
Prowl H2O <sup>g</sup>	_	4	_	_	_	_	_	_	_	_	4	_	_	_	_	_	
Pummel	_	_			_	-	_	_	_	_	4.5	_	_	_	_	_	
Pursuit <sup>h</sup>	_	_	4	_	0	0	_	_	_	_	4	_	4	_	_	_	
Python	-	4	4		4	4	—	_	—	_	4	-	_	4	—	-	
Rowel <sup>i</sup>	4	3	3	4	3	3	4	1	_	4	1	_	4	4	4	4	
Rowel FX	4	4	4	_	—	-	_	_	_	_	4	-	_	_	-	-	
Sharpen <sup>j</sup>	4	0	0	2	4	2	0	0	4	4	4	4	4	0	0	4	
Sonic	_	_	_	_	_	_	_	_	_	_	4	_	_	_	_	_	
Surveil Co Pack	-	_			—	-	—	_	—	_	3	-	_	_	—	-	
Torment	_	_	4	_	_	_	4	4	_	_	4	_	_	4	4	_	
Trivence	_	4	_	-	—	_	—	—	—	—	4	—	—	_	—	—	
Valor SX <sup>k</sup>	_	3-4	3-4	-	3-4	3-4	_	1-9	_	_	1-9	_	4-10	4-10	_	_	
Valor XLT <sup>I</sup>	_	4	4	-	—	_	—	—	—	—	4	—	—	_	—	—	
Vise	_	4.5	4.5		_	-	_	_	_	—	4.5	-	_	4.5	—	-	
Warrant	_	_	_	-	—	_	—	—	—	—	4	—	—	_	—	—	
Warrant Ultra	_	4	4	-	—	_	—	—	—	—	4	—	—	4	—	—	

See footnotes on next page.

<sup>a</sup> Wheat may be planted when Anthem is applied at less than 9.75 oz/ac. Field pea may be planted when Anthem is applied at less than 3.25 oz/ac.

<sup>b</sup> Wheat may be planted when Anthem Maxx is applied at less than 4.875 oz/ac. Field pea may be planted when Anthem Maxx is applied at less than 1.625 oz/ac.

<sup>c</sup> Intervals for cereal rye and wheat are 18 months for Minnesota and North Dakota, north of Highway 210. Do NOT feed treated soybean forage or hay to livestock.

<sup>d</sup> Do not graze or harvest for food or feed cover crops planted less than nine months after treatment with Command.

<sup>e</sup> Peas may be planted immediately following application when Command is applied at less than 0.5 lb/ac; otherwise, the interval is nine months.

<sup>f</sup> Intervals are for rates of Dual II Magnum applied at NOT more than 1.33 pint/ac. A second application or other herbicides containing *S*-metolachlor is prohibited when planting forage double crops in the fall.

<sup>g</sup> When applied at a rate of 3.2 pts/ac.

<sup>h</sup> This will restrict the post-emergence application of Pursuit in soybean if growers intend to plant alfalfa, cereal rye, clover, or wheat in the fall.

<sup>i</sup> A rotational interval for most cover crops is four months if soil is tilled prior to planting, and eight months if no tillage is performed. See herbicide label for instructions.

<sup>j</sup> Can be applied at 1 fl oz/ac if the organic matter content of the soil is less than 2%. This herbicide will provide limited broadleaf control at this rate. Coarse soils with less than 2% organic matter require a minimum preplant interval of 30 days for planting soybean.

<sup>k</sup> Rotation intervals for Valor SX vary depending on the applied rate of herbicide; see label for full restrictions.

<sup>1</sup> Valor XLT in Nebraska can only be applied in soybean fields south of Route 30 and east of Route 281.