

**Table 2. Corn burndown and PRE herbicides that will not cause carryover injury in field peas.**

Herbicide table includes rotation restrictions for field peas and efficacy rating for NE most troublesome weeds

Site of Action	Corn herbicide	Common name	Rotation Restriction for Field Peas <sup>1</sup>	Weed Response to Selected Herbicides					
				Kochia	Lambsquarters	Marestail	Pigweeds	Russian Thistle	Foxtail species
<b>Burndown</b>									
4	2,4-D ester 4L	2,4-D	NCS	7	9	7	-	9	1
14	Aim	carfentrazone-ethyl	AT	6	9	-	7-8	6	1
14+15	Anthem MAXX	pyroxasulfone + fluthiacet-methyl	6	7	9	8	-	9	1
4	Banvel/dicamba	Dimethylamine salt of dicamba	NCS <sup>g</sup>	9	9	8	-	5	1
2	Basis Blend	rimsulfuron + thifensulfuron	8/10 <sup>ab</sup>	3	7	5	9	4	7
4	Clarity	dicamba-glycolamine	180 D <sup>m</sup>	9	9	8	-	5	1
4+27	Diflex DUO	dicamba-glycolamine + tembotrione	10	9	9	8	-	5	2
19+4	Distinct	diflufenopyr + dicamba	4	9	7	9	-	9	1
14+15	Fierce	flumioxazin + pyroxasulfone	6/11 <sup>dq</sup>		check 2018 weed guide for tank mixes				
9	Glyphosate	glyphosate	AT	8	8	8	-	9	9
22	Gramoxone SL	paraquat	AT	9	7	7	-	6	7
9+4	Land Master BW	glyphosate + 2,4-D amine	3	9	9	9	-	9	9
4	Python	flumetsulam	4		check 2018 weed guide for tank mixes				
14	Sharpen + glyphosate	saflufenacil	4-9 <sup>cl</sup>	9	9	10	-	10	10
14	Valor SX	flumioxazin + chlorimuron ethyl	4		check 2018 weed guide for tank mixes				
14+15	Verdict + glyphosate	saflufenacil + dimethenamid-P	NCS	9	8	9	-	10	10
<b>PRE</b>									
14+15	Anthem MAXX	pyroxasulfone + fluthiacet-methyl	6	6	7	-	9	5	9
15	Breakfree NXT/Harness	acetochlor	NCS	2	7	-	7	3	9
15	Surpass NXT/Confidence	acetochlor	NCS	2	7	-	7	3	9
15	Dual/Chich/Parallel	S-metolachlor + benoxacor	AT	2	7	-	7	3	9
15	Degree/Topnotch	acetochlor + safener	NCS	2	8	-	8	5	9
4+2	Hornet WDG	flumetsulam + clopyralid	10.5/18 <sup>p</sup>	9	9	-	9	8	1
15	Outlook	dimethenamid-P	NCS	2	7		8	3	9
3	Prowl H2O	pentimethalin	NCS		check 2018 weed guide for tank mixes				
2	Resolve SG	rimsulfuron	10	8	7	-	9	3	8
14	Sharpen	saflufenacil	4-9 <sup>cl</sup>	7	8	-	9	8	2
15+4+2	SureStart II/Triple Flex	acetochlor + clopyralid + flumetsulam	NCS						
14+15	Verdict	saflufenacil + dimethenamid-P	NCS	7	9	-	10	9	9
15	Zidua	pyroxasulfone	6	7	7	-	8	6	8

<sup>1</sup> Months unless otherwise noted. D = Days; AT = Any Time; NCS = Next Cropping Season; 2CS = Second Cropping Season; 3CS Tird Cropping Season, NTE = No Tolerance Established NI = No information, FBA = Field Bioassay, DNR = Do not rotate.

<sup>ab</sup> 11 months for STS varieties, 36 months or earlier with a bioassay; 14 or 26 months if pH < 7.9 and rainfall limits are followed.

<sup>cl</sup> Rotation interval depends upon rate applied and soil texture. See the label for detailed instructions

<sup>dq</sup> 6 months for field peas, 11 month for edible peas

<sup>g</sup> Rotation interval is 45 days per pint of Banvel applied at 23 days per pint of Banvel SGF, excluding days when the ground is frozen

<sup>m</sup> Applications of 24 oz/A or less = 22 days for each 8 fluid oz; 24 oz/A or more = 45 day interval for each 16 fluid oz/A applied.