

Management of Corn Diseases

Fungicide Efficacy for Control of Corn Diseases—April 2014

The Corn Disease Working Group (CDWG) has developed the following information on fungicide efficacy for control of major corn diseases in the United States. Efficacy ratings for each fungicide listed in the table were determined by field testing the materials over multiple years and locations by the members of the committee. Efficacy ratings are based upon level of disease control achieved by product, and are not necessarily reflective of yield increases obtained from product application. Efficacy depends upon proper application timing, rate, and application method to achieve optimum effectiveness of the fungicide as determined by labeled instructions and overall level of disease in the field at the time of application. Differences in efficacy among fungicide products were determined by direct comparisons among products in field tests and are based on a *single application* of the labeled rate as listed in the table. **Table includes systemic fungicides available that have been tested over multiple years and locations. The table is not intended to be a list of all labeled products¹.** Efficacy categories: NR=Not Recommended; P=Poor; F=Fair; G=Good; VG=Very Good; E=Excellent; NL = Not Labeled for use against this disease; U = Unknown efficacy or insufficient data to rank product

Fungicide(s)				Anthracnose leaf blight	Common rust	Eyespot	Gray leaf spot	Northern leaf blight	Southern rust	Harvest Restriction ²
Class	Active ingredient (%)	Product/Trade name	Rate/A (fl oz)							
QoI Strobilurins Group 11	Azoxystrobin 22.9%	Quadris 2.08 SC	6.0 - 15.5	VG	E	VG	E	G	G	7 days
	Pyraclostrobin 23.6%	Headline 2.09 EC/SC	6.0 - 12.0	VG	E	E	E	VG	E	7 days
	Picoxystrobin	Approach 2.08 SC	3.0 – 12.0	VG	VG-E	VG	F-VG	VG	U	7 days
DMI Triazoles Group 3	Propiconazole 41.8%	Tilt 3.6 EC Multiple Generics	2.0 - 4.0	NL	VG	E	G	G	G	30 days
	Prothioconazole 41.0%	Proline 480 SC	5.7	U	VG	E	U	VG	G	14 days
	Tebuconazole 38.7%	Folicur 3.6 F Multiple Generics	4.0 - 6.0	NL	U	NL	U	VG	U	36 days
	Tetraconazole 20.5%	Domark 230 ME	4.0 – 6.0	U	U	U	E	U	G	R3 (milk)
Mixed mode of action	Azoxystrobin 7.0% Propiconazole 11.7%	Quilt 200 SC	7.0 - 14.0	U	VG-E	E	E	VG	VG	30 days
	Azoxystrobin 13.5% Propiconazole 11.7%	Quilt Xcel 2.2 SE	10.5 - 14.0	VG	VG-E	VG-E	E	VG	VG	30 days
	Pyraclostrobin 13.6% Metconazole 5.1%	Headline AMP 1.68 SC	10.0 - 14.4	U	E	E	E	VG	VG	20 days
	Pyraclostrobin 28.58% Fluxapyroxad 14.33%	Priaxor 4.17 SC	4.0 – 8.0	U	VG	U	VG	U	G	21 days
	Trifloxystrobin 11.4% Propiconazole 11.4%	Stratego 250 EC	10.0 - 12.0	U	VG	E	VG	G	G	30 days
	Trifloxystrobin 32.3% Prothioconazole 10.8%	Stratego YLD 4.18 SC	4.0 - 5.0	VG	E	VG	E	VG	VG	14 days

¹Additional fungicides are labeled for disease on corn, including contact fungicides such as chlorothalonil. Certain fungicides may be available for diseases not listed in the table, including Gibberella and Fusarium ear rot. Applications of Proline 480 SC for use on ear rots requires a FIFRA Section 2(ee) and is only approved for use in Illinois, Indiana, Iowa, Louisiana, Maryland, Michigan, Mississippi, North Dakota, Ohio, Pennsylvania, and Virginia.

²Harvest restrictions are listed for field corn harvested for grain. Restrictions may vary for other types of corn (sweet, seed or popcorn, etc.), and corn for other uses such as forage or fodder.