

**Table 7. Field peas – Rotation restrictions for selected field pea herbicides**

Site of Action	Field peas herbicide	Rotation Restrictions for selected crops																				
		Grass crops												Broadleaf crops								
		Field Corn	Seed Corn	Popcorn	Sweet Corn	Winter Wheat	Spring Wheat	Oat	Winter Barley	Spring Barley	Rye	Grain Sorghum	Proso Millet	Soybean	Canola	Buckwheat	Sunflower	Sugarbeet	Dry Bean	Potato	Alfalfa	Red Clover
	<b>PRE</b>																					
14+15	BroadAxe XC/SpartanElite	10	10	18	18	4.5	4.5	12	4.5	4.5	4.5	10	12	AT	NTE	12	AT	36	NTE	4	12	NTE
15	Dual II Magnum	AT	AT	AT	AT	4.5	4.5	4.5	4.5	NCS	3	7D/14D <sup>cs</sup>	NCS	15D/30D <sup>cs</sup>	NCS	NCS	NCS	NCS	NCS	NCS	NCS	NCS
14+2	Optil	8.5 <sup>ck</sup>	8.5	18	18	4 <sup>ck</sup>	4 <sup>ck</sup>	18	9.5	9.5	4-18	18	40	0-1	9.5 <sup>ck</sup>	40	9.5-18 <sup>ck</sup>	40	4	26	4	4
3	Prowl H2O	NCS	NCS	NCS	NCS	4 <sup>**</sup>	NCS	NCS	4 <sup>**</sup>	NCS	NCS	NCS	NCS	NCS	NCS	NCS	NCS	NCS	NCS	NCS	NCS	NCS
14	Sharpen	AT	AT	AT	0.5	0-3 <sup>cl</sup>	0-3 <sup>cl</sup>	0-3 <sup>cl</sup>	0-3 <sup>cl</sup>	0-3 <sup>cl</sup>	0-3 <sup>cl</sup>	0-1 <sup>cl</sup>	0-6 <sup>cl</sup>	4-9 <sup>cl</sup>	4-9 <sup>cl</sup>	4-9 <sup>cl</sup>	4-9 <sup>cl</sup>	4-9 <sup>cl</sup>	4-9 <sup>cl</sup>	4-9 <sup>cl</sup>	4-9 <sup>cl</sup>	4-9 <sup>cl</sup>
14	Spartan	10	12	12	18	18	4	4	12	4	4	10/18 <sup>by</sup>	12	AT	24	12/FBA	AT	36	12	12/FBA	12	12
14	Spartan Charge	4	12	12	12	4	4	12	4	3	3	10/18 <sup>by</sup>	12	AT	10/18 <sup>v</sup>	18	18	18	10/18 <sup>v</sup>	10/18 <sup>v</sup>	18	18
3	Treflan (PPI)	NCS	NCS	NCS	NCS	18	2CS	12/18 <sup>ba</sup>	2CS	2CS	2CS	12/18 <sup>ba</sup>	NCS	AT	NCS	NCS	AT	12/14 <sup>az</sup>	AT	AT	NCS	NCS
	<b>POST</b>																					
1	Assure II	4	4	4	4	4	4	4	4	4	4	4	4	AT	AT	4	4	AT	AT	4	4	4
6	Basagran 5L	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT	AT
1	Poast	4 <sup>bf</sup>	4	4	4	4	4	4	4	4	4	4	4	AT	4	4	AT	AT	AT	AT	AT	AT
2	Pursuit	8.5 <sup>h</sup>	8.5	18	18	4	4	18	9.5	4	4	12	26/FBA	AT	26/FBA	26/FBA	18	26/FBA	4	12	4	26/FBA
6	Raptor	8.5	8.5	8.5	8.5	0/3 <sup>bv</sup>	0/3 <sup>bv</sup>	9	4 <sup>ca</sup>	4 <sup>ca</sup>	3	9	9	AT	18-26	18	9	18-26 <sup>as,bv</sup>	AT	18	3	18
1	Select Max	6 D	1	1	1	1	1	1	1	1	1	1	1	AT	1	1	1	AT	AT	1	AT	1
2+6	Varisto <sup>a</sup>	8.5	8.5	8.5	8.5	3	3	9	9	9	4	9	9	AT	26	18	9	26	AT	18	3	NTE

<sup>1</sup> Months unless otherwise noted. D = Days; AT = Any Time; NCS = Next Cropping Season; 2CS = Second Cropping Season; 3CS Tird Cropping Season, NTE = No Tolerance Established, NI = No information, FBA = Field Bioassay, DNR = Do not rotate.  
<sup>as</sup> 18 months in eastern Nebraska if soil pH is 6.2 or grater and 26 months if the soil pH is less than 6.2; 26 months for western Nebraska  
<sup>ba</sup> All areas receiving more than 20 inches of rainfall and irrigation - those areas receiving less than 20 inches of rainfall and irrigation to produce a crop.  
<sup>bf</sup> Poast protected field corn hybrids may be planted anytime  
<sup>bv</sup> Clearfield/normal non-Clearfield)  
<sup>by</sup> 18-month rotation for rates above 0.25 lb ai/A sulfentrazone.  
<sup>ca</sup> Applied prior to June 1 in previous year.  
<sup>ck</sup> Check the label for specific rotations for Clearfield crops (corn, wheat, canola, and sunflower).  
<sup>cl</sup> Rotation interval depends upon rate applied and soil texture. See the label for detailed instructions  
<sup>cs</sup> Rotation interval should be extended to 18 months if drought conditions prevail after application unless at least 15 inch of sprinkler irrigation has been applied  
<sup>h</sup> Clearfield, IR, or IMR field corn hybrids may be planted "anytime".  
<sup>v</sup> Rotation interval varies by location in Nebraska, soil pH, application rate, and cummulative precipitation